

incompatible incompatibilities

and how to make them compatible again

Francesco Buscemi, Nagoya University

Quantum TUT Workshop 2024
Toyohashi University of Technology
Toyohashi, 22 February 2024



references

- F.B., E. Chitambar, W. Zhou:
A complete resource theory of quantum (POVMs) incompatibility as quantum programmability.
Physical Review Letters 124, 120401 (2020)
- F.B., K. Kobayashi, S. Minagawa, P. Perinotti, A. Tosini:
Unifying different notions of quantum (instruments) incompatibility into a strict hierarchy of resource theories of communication.
Quantum 7, 1035 (2023)

POVMs and instruments

in this talk: all sets (\mathbb{X}, \mathbb{Y} etc.) are finite, all spaces ($\mathcal{H}_A, \mathcal{H}_B$ etc.) are finite-dimensional

POVM: family \mathbf{P} of positive semidefinite operators on \mathcal{H} labeled by set \mathbb{X} , i.e., $\mathbf{P} = \{P_x\}_{x \in \mathbb{X}}$, with $P_x \geq 0$ and $\sum_x P_x = \mathbb{1}$

interpretation: expected probability of outcome x is $p(x) = \text{Tr}[\rho P_x]$

POVMs and instruments

in this talk: all sets (\mathbb{X}, \mathbb{Y} etc.) are finite, all spaces ($\mathcal{H}_A, \mathcal{H}_B$ etc.) are finite-dimensional

POVM: family \mathbf{P} of positive semidefinite operators on \mathcal{H} labeled by set \mathbb{X} , i.e., $\mathbf{P} = \{P_x\}_{x \in \mathbb{X}}$, with $P_x \geq 0$ and $\sum_x P_x = \mathbb{1}$

interpretation: expected probability of outcome x is $p(x) = \text{Tr}[\varrho P_x]$

instrument: family $\{\mathcal{I}_x : A \rightarrow B\}_{x \in \mathbb{X}}$ of completely positive (CP) linear maps from $\mathcal{B}(\mathcal{H}_A)$ to $\mathcal{B}(\mathcal{H}_B)$, such that $\sum_x \mathcal{I}_x$ is trace-preserving (TP)

interpretation: expected probability of outcome x is $p(x) = \text{Tr}[\mathcal{I}_x(\varrho)]$, and corresponding post-measurement state is $\frac{1}{p(x)} \mathcal{I}_x(\varrho)$

incompatibility

In quantum theory, some measurements necessarily exclude others.

If all measurements were compatible, we would not have QKD, violation of Bell's inequalities, quantum speedups, etc.

Various formalizations:

- preparation uncertainty relations
- measurement (e.g., error-disturbance) uncertainty relations
- incompatibility

compatible POVMs 1/2

Definition

given a family $\{\mathbf{P}^{(i)}\}_{i \in \mathbb{I}} \equiv \{P_x^{(i)}\}_{x \in \mathbb{X}, i \in \mathbb{I}}$ of POVMs, all defined on the same system A , we say that the family is **compatible**, whenever there exists

- a “mother” POVM $\mathbf{O} = \{O_w\}_{w \in \mathbb{W}}$ on system A
- a conditional probability distribution $\mu(x|w, i)$

such that

$$P_x^{(i)} = \sum_w \mu(x|w, i) O_w ,$$

for all $x \in \mathbb{X}$ and all $i \in \mathbb{I}$.


But what does it mean, *operationally*, if I say that, e.g., a certain laboratory can only perform compatible measurement?

compatible POVMs 2/2

There's a bipartition hidden in the concept of (in)compatibility:

compatible POVMs 2/2

There's a bipartition hidden in the concept of (in)compatibility:



[F.B., E. Chitambar, W. Zhou; PRL 2020]

the problem

While there is consensus on a single notion of **compatibility** for POVMs, in the case of **instruments**, the situation is less clear...

classical compatibility 1/2

Definition (Heinosaari–Miyadera–Reitzner, 2014)

given a family of instruments $\{\mathcal{I}_x^{(i)} : A \rightarrow B\}_{x \in \mathbb{X}, i \in \mathbb{I}}$, we say that the family is **classically compatible**, whenever there exist


- a **mother instrument** $\{\mathcal{H}_w : A \rightarrow B\}_{w \in \mathbb{W}}$
- a **conditional probability distribution** $\mu(x|w, i)$

such that

$$\mathcal{I}_x^{(i)} = \sum_w \mu(x|w, i) \mathcal{H}_w ,$$

for all $x \in \mathbb{X}$ and all $i \in \mathbb{I}$.

classical compatibility 2/2



crucially:

- **II is classical**: no shared entanglement, communication is classical
- **the box is II \rightarrow I non-signaling**: communication goes only from **I** to **II**; see [Ji and Chitambar; PRA 2021]

alternative: marginalizing the mother

alternative: marginalizing the mother

- in the case of POVMs: without loss of generality (because classical information can be copied), one can consider only **marginalizations**, i.e.,

$$P_x^{(i)} = \sum_{x_j: j \neq i} O_{x_1, x_2, \dots, x_n}$$

alternative: marginalizing the mother

- in the case of POVMs: without loss of generality (because classical information can be copied), one can consider only **marginalizations**, i.e.,

$$P_x^{(i)} = \sum_{x_j: j \neq i} O_{x_1, x_2, \dots, x_n}$$

- the notion of “**parallel compatibility**” for instruments applies the same intuition to the quantum outputs too

parallel compatibility 1/2

Definition (Heinosaari–Miyadera–Ziman, 2015)

given a family of instruments $\{\mathcal{I}_x^{(i)} : A \rightarrow B_i\}_{x \in \mathbb{X}, i \in \mathbb{I}}$, we say that the family is **parallelly compatible**, whenever there exist


- a **mother instrument** $\{\mathcal{H}_w : A \rightarrow \bigotimes_{i \in \mathbb{I}} B_i\}_{w \in \mathbb{W}}$
- a **conditional probability distribution** $\mu(x|w, i)$

such that


$$\mathcal{I}_x^{(i)} = \sum_w \mu(x|w, i) [\text{Tr}_{B_i^c} \circ \mathcal{H}_w], \quad B_i^c := \bigotimes_{i' \in \mathbb{I}: i' \neq i} B_{i'},$$

for all $x \in \mathbb{X}$ and all $i \in \mathbb{I}$.

parallel compatibility 2/2




parallel compatibility VS classical compatibility




- parallel compatibility is both $\text{I} \rightarrow \text{II}$ and $\text{II} \rightarrow \text{I}$ signaling; therefore, **parallel compatibility $\not\Rightarrow$ classical compatibility**

parallel compatibility VS classical compatibility



- parallel compatibility is both $\text{I} \rightarrow \text{II}$ and $\text{II} \rightarrow \text{I}$ signaling; therefore, **parallel compatibility** $\not\Rightarrow$ **classical compatibility**
- non-disturbing instruments are never parallelly compatible; therefore, **classical compatibility** $\not\Rightarrow$ **parallel compatibility**

parallel compatibility VS classical compatibility



- parallel compatibility is both $\text{I} \rightarrow \text{II}$ and $\text{II} \rightarrow \text{I}$ signaling; therefore, **parallel compatibility** $\not\Rightarrow$ **classical compatibility**
- non-disturbing instruments are never parallelly compatible; therefore, **classical compatibility** $\not\Rightarrow$ **parallel compatibility**
- parallel compatibility is more closely related to **quantum no-broadcasting** than it is to measurement compatibility

bridging the two camps

q-compatibility

Definition

given a family of instruments $\{\mathcal{I}_x^{(i)} : A \rightarrow B_i\}_{x \in \mathbb{X}, i \in \mathbb{I}}$, we say that the family is **q-compatible**, whenever there exist


- a **mother instrument** $\{\mathcal{H}_w : A \rightarrow C\}_{w \in \mathbb{W}}$
- a **conditional probability distribution** $\mu(x|w, i)$
- a **family of postprocessing channels** $\{\mathcal{D}^{(x,w,i)} : C \rightarrow B_i\}_{x \in \mathbb{X}, w \in \mathbb{W}, i \in \mathbb{I}}$

such that

$$\mathcal{I}_x^{(i)} = \sum_w \mu(x|w, i) [\mathcal{D}^{(x,w,i)} \circ \mathcal{H}_w] ,$$

for all $x \in \mathbb{X}$ and all $i \in \mathbb{I}$.

q-compatibility as a circuit




Special cases:


- **classical compatibility**: when $C \equiv B_i$ for all i and $\mathcal{D}^{(x,w,i)} = \text{id}$
- **parallel compatibility**: when $C \equiv \bigotimes_i B_i$ and $\mathcal{D}^{(x,w,i)} = \text{Tr}_{B_i^c}$

incompatibility-non-increasing operations

free operations for classical incompatibility




free operations for classical incompatibility




- all classically compatible devices can be created for free
- if the initial device (the dark gray inner box) is classically compatible, the final device is also classically compatible

free operations for q-incompatibility



free operations for q-incompatibility



- all q-compatible devices can be created for free
- if the initial device (the dark gray inner box) is q-compatible, the final device is also q-compatible

beyond q-compatibility

simultaneous VS sequential compatibility

compatibility of POVMs is always, w.l.o.g., **simultaneous compatibility** (again, because classical information can be copied)

simultaneous VS sequential compatibility

compatibility of POVMs is always, w.l.o.g., **simultaneous compatibility** (again, because classical information can be copied)

for instruments, it is more subtle: for example, consider the following two instruments

$$\mathcal{I}_1(\bullet) = pU_1 \bullet U_1^\dagger$$

$$\mathcal{I}_2(\bullet) = (1 - p)U_2 \bullet U_2^\dagger$$

$$\mathcal{J}_1(\bullet) = |0\rangle\langle 0| \bullet |0\rangle\langle 0|$$

$$\mathcal{J}_2(\bullet) = |1\rangle\langle 1| \bullet |1\rangle\langle 1|$$

the corresponding POVMs, i.e., $\{p\mathbb{1}, (1 - p)\mathbb{1}\}$ and $\{|0\rangle\langle 0|, |1\rangle\langle 1|\}$ are compatible, but **the two instruments are not** (they are not even q-compatible)

simultaneous VS sequential compatibility

compatibility of POVMs is always, w.l.o.g., **simultaneous compatibility** (again, because classical information can be copied)

for instruments, it is more subtle: for example, consider the following two instruments

$$\mathcal{I}_1(\bullet) = pU_1 \bullet U_1^\dagger$$

$$\mathcal{I}_2(\bullet) = (1-p)U_2 \bullet U_2^\dagger$$

$$\mathcal{J}_1(\bullet) = |0\rangle\langle 0| \bullet |0\rangle\langle 0|$$

$$\mathcal{J}_2(\bullet) = |1\rangle\langle 1| \bullet |1\rangle\langle 1|$$

the corresponding POVMs, i.e., $\{p\mathbb{1}, (1-p)\mathbb{1}\}$ and $\{|0\rangle\langle 0|, |1\rangle\langle 1|\}$ are compatible, but **the two instruments are not** (they are not even q-compatible)

except that, in a sense, they actually *are* compatible!

just not simultaneously so: do \mathcal{I} , keep the outcome, undo the unitary, and finally do \mathcal{J}

simultaneous VS sequential compatibility

compatibility of POVMs is always, w.l.o.g., **simultaneous compatibility** (again, because classical information can be copied)

for instruments, it is more subtle: for example, consider the following two instruments

$$\begin{aligned}\mathcal{I}_1(\bullet) &= pU_1 \bullet U_1^\dagger & \mathcal{J}_1(\bullet) &= |0\rangle\langle 0| \bullet |0\rangle\langle 0| \\ \mathcal{I}_2(\bullet) &= (1-p)U_2 \bullet U_2^\dagger & \mathcal{J}_2(\bullet) &= |1\rangle\langle 1| \bullet |1\rangle\langle 1|\end{aligned}$$

the corresponding POVMs, i.e., $\{p\mathbb{1}, (1-p)\mathbb{1}\}$ and $\{|0\rangle\langle 0|, |1\rangle\langle 1|\}$ are compatible, but **the two instruments are not** (they are not even q-compatible)

except that, in a sense, they actually *are* compatible!

just not simultaneously so: do \mathcal{I} , keep the outcome, undo the unitary, and finally do \mathcal{J}

remark: reversing the order (i.e., first \mathcal{J} , then \mathcal{I}) the same construction does not work

no-exclusivity

Definition

an instrument $\{\mathcal{I}_x : A \rightarrow B_1\}_{x \in \mathbb{X}}$ **does not exclude** another instrument $\{\mathcal{J}_y : A \rightarrow B_2\}_{y \in \mathbb{Y}}$, whenever there exist


- a **mother instrument** $\{\mathcal{H}_w : A \rightarrow C\}_{w \in \mathbb{W}}$
- a **conditional probability distribution** $\mu(x|w)$
- a **family of postprocessing channels** $\{\mathcal{D}^{(x,w)} : C \rightarrow B_1\}_{x \in \mathbb{X}, w \in \mathbb{W}}$
- a **family of instruments** $\{\mathcal{K}_y^{(w)} : C \rightarrow B_2\}_{w \in \mathbb{W}, y \in \mathbb{Y}}$

such that


$$\mathcal{I}_x = \sum_w \mu(x|w) [\mathcal{D}^{(x,w)} \circ \mathcal{H}_w], \quad \mathcal{J}_y = \sum_w \mathcal{K}_y^{(w)} \circ \mathcal{H}_w,$$

for all $x \in \mathbb{X}$ and all $y \in \mathbb{Y}$.

no-exclusivity as a circuit




no-exclusivity VS q-compatibility



in the definition of no-exclusivity, the post-processing box at I can be an instrument

no-exclusivity VS q-compatibility



in the definition of no-exclusivity, the post-processing box at I can be an instrument

intuition: the information necessary to reconstruct the result of one instrument (i.e., the non-excluding one) can be extracted with a disturbance “small enough” to not exclude the other instrument (i.e., the non-excluded one)

formulating resource theories of incompatibility

a hierarchy of incompatibility preorders

given two families of instruments $\{\mathcal{I}_x^{(i)} : A \rightarrow B_i\}_{x \in \mathbb{X}, i \in \mathbb{I}}$ and $\{\mathcal{J}_y^{(j)} : C \rightarrow D_j\}_{y \in \mathbb{Y}, j \in \mathbb{J}}$,
we say

$\{\mathcal{I}_x^{(i)} : A \rightarrow B_i\}$ is **more incompatible/exclusive** than $\{\mathcal{J}_y^{(j)} : C \rightarrow D_j\}$

whenever the former can be transformed into the latter by means of a corresponding
free operation

a hierarchy of incompatibility preorders

given two families of instruments $\{\mathcal{I}_x^{(i)} : A \rightarrow B_i\}_{x \in \mathbb{X}, i \in \mathbb{I}}$ and $\{\mathcal{J}_y^{(j)} : C \rightarrow D_j\}_{y \in \mathbb{Y}, j \in \mathbb{J}}$,
we say

$\{\mathcal{I}_x^{(i)} : A \rightarrow B_i\}$ is **more incompatible/exclusive** than $\{\mathcal{J}_y^{(j)} : C \rightarrow D_j\}$

whenever the former can be transformed into the latter by means of a corresponding
free operation

more classically incompatible \implies more q-incompatible \implies more exclusive

a hierarchy of incompatibility preorders

given two families of instruments $\{\mathcal{I}_x^{(i)} : A \rightarrow B_i\}_{x \in \mathbb{X}, i \in \mathbb{I}}$ and $\{\mathcal{J}_y^{(j)} : C \rightarrow D_j\}_{y \in \mathbb{Y}, j \in \mathbb{J}}$, we say

$\{\mathcal{I}_x^{(i)} : A \rightarrow B_i\}$ is **more incompatible/exclusive** than $\{\mathcal{J}_y^{(j)} : C \rightarrow D_j\}$

whenever the former can be transformed into the latter by means of a corresponding free operation

more classically incompatible \implies more q-incompatible \implies more exclusive

\rightsquigarrow this is now an instance of *statistical comparison*: a complete family of monotones can be constructed and a **Blackwell-like theorem proved**

further details on arXiv:2211.09226

conclusion

take home messages

- no need to argue about the “correct” definition of compatibility for instruments:
q-compatibility provides an overarching framework

take home messages

- no need to argue about the “correct” definition of compatibility for instruments: **q-compatibility provides an overarching framework**
- as such, we can think of it as a **resource**, and construct a whole hierarchy of complete resource theories of incompatibility, all of which fall back on the same notion of compatibility of POVMs in the case of instruments with trivial quantum output

take home messages

- no need to argue about the “correct” definition of compatibility for instruments: **q-compatibility provides an overarching framework**
- as such, we can think of it as a **resource**, and construct a whole hierarchy of complete resource theories of incompatibility, all of which fall back on the same notion of compatibility of POVMs in the case of instruments with trivial quantum output
- for instruments, simultaneous compatibility is not the end of the story: we also have a notion of sequential compatibility, i.e., **no-exclusivity**

take home messages

- no need to argue about the “correct” definition of compatibility for instruments: **q-compatibility provides an overarching framework**
- as such, we can think of it as a **resource**, and construct a whole hierarchy of complete resource theories of incompatibility, all of which fall back on the same notion of compatibility of POVMs in the case of instruments with trivial quantum output
- for instruments, simultaneous compatibility is not the end of the story: we also have a notion of sequential compatibility, i.e., **no-exclusivity**

thank you for your attention